

IMMUNEX CORPORATION

Attorney Docket No. 2623-A

TITLE

METHOD OF REGULATING NITRIC OXIDE PRODUCTION

CROSS-REFERENCE TO RELATED APPLICATION

5 This application is a continuation of U.S. Application Serial Number 07/507,213,
 filed November 27, 1996.

TECHNICAL FIELD OF THE INVENTION

10 The present invention relates generally to the modulation of levels of nitric oxide,
 particularly in osteoarthritis.

BACKGROUND OF THE INVENTION

15 Cytokines are hormone-like molecules that regulate various aspects of an immune
 or inflammatory response; they exert their effects by specifically binding receptors present
 on cells, and transducing a signal to the cells. In addition to having beneficial effects (i.e.,
 development of an effective immune response and control of infectious disease), cytokines
 have also been implicated in various autoimmune and inflammatory conditions.

20 Various cartilage associated cells (i.e., chondrocytes, synovial lining cells,
 endothelial cells, synovial fibroblasts and mononuclear cells that are present in a joint) can
 release nitric oxide (NO). This free radical serves as a front-line antimicrobial agent and
 also has antitumor effects. However, NO has also been implicated in several deleterious
 conditions, including autoimmune and inflammatory diseases and the bone destruction that
 occurs in osteoarthritis, which is not typically thought of as an inflammatory condition.

25 Rouvier et al. (*J. Immunol.* 150:5445; 1993) reported a novel cDNA which they
 termed CTLA-8, and which has since become known as Interleukin-17 (IL-17). IL-17 is
 57% homologous to the predicted amino acid sequence of an open reading frame (ORF)
 present in Herpesvirus saimiri (HSV) referred to as HVS13 (Nicholas et al. *Virol.* 179:1
 89, 1990; Albrecht et al., *J. Virol.* 66:5047; 1992).

30 A novel receptor that binds IL-17 and its viral homolog, HVS13, has been cloned
 as described in USSN 08/620,694, filed March 21, 1996. The receptor is a Type I
 transmembrane protein; the mouse receptor has 864 amino acid residues, the human
 receptor has 866 amino acid residues. A soluble form of the receptor was found to inhibit
 various IL-17-mediated activities.

SUMMARY OF THE INVENTION

35 Nitric oxide (NO) is a free radical that is involved in many phenomena, including
 the pathophysiological conditions of rheumatoid arthritis (RA) and osteoarthritis (OA). IL-
 17 stimulates production of NO by cartilage from individuals afflicted with OA. A soluble

form of IL-17R was found to inhibit various IL-17-mediated activities. Accordingly, soluble IL-17R will be useful in regulating levels of NO in a clinical setting.

DETAILED DESCRIPTION OF THE INVENTION

5 Nitric oxide is an intracellular signaling molecule that is involved in many physiological phenomena, including endothelium-dependent relaxation, neurotransmission and cell-mediated immune responses. As an antimicrobial agent, NO is effective against bacteria, viruses, helminths and parasites; it is also useful in the killing of tumor cells. Increased levels of NO occur in inflammatory disease (i.e., arthritis, ulcerative colitis,
10 diabetes, Crohn's disease), and inhibitors of NO synthetases (NOS) have been used in experimental models of inflammatory disease, with varied effects (reviewed by A.O. Vladutiu in *Clinical Immunology and Immunopathology* 76:1-11; 1995).

Osteoarthritis (OA) has typically been considered a non-inflammatory disease, however, Amin et al. (*J. Exp. Med.* 182:2097; 1995) recently reported that the levels of
15 NOS are upregulated in cartilage from OA patients. Incubation of OA-affected cartilage in serum-free medium resulted in the spontaneous release of substantial amounts of NO. Interleukin-18 (IL-18), tumor necrosis factor- α (TNF- α) and lipopolysaccharide (LPS) augmented the nitrite release of OA-affected cartilage. Similar results were observed by Sakurai et al. (*J. Clin. Invest.* 96:2357, 1995) for rheumatoid arthritis patients.

20 IL-17 also upregulates release of NO from OA-affected cartilage. Moreover, inhibitors of IL-18 and TNF- α do not inhibit the IL-17-augmented release of NO. Accordingly, inhibitors of IL-17 will be useful in regulating levels of NO. Such inhibitors will find therapeutic application in ameliorating the effects of NO in OA, as well as in other disease conditions in which this free radical plays a role (i.e., autoimmune and
25 inflammatory disease).

A particularly preferred form of IL-17 inhibitor is soluble IL-17R, which is described in detail in USSN 08/620,694. IL-17 inhibitors may be used in conjunction with (i.e., simultaneously, separately or sequentially) inhibitors of IL-1 and TNF. Exemplary
30 IL-1 inhibitors include soluble IL-1 receptors such as those described in U.S. Patents 5,319,071, 5,180,812 and 5,350,683, as well as a protein known as IL-1 receptor antagonist (IL-1RA; Eisenberg et al., *Nature* 343:341, 1990) and inhibitors of an enzyme that cleaves IL-1 into its biologically active form, as described in U.S. Patent 5,416,013.

Exemplary TNF inhibitors include soluble forms of TNF receptors, for example as described in U.S. Patent 5,395,760, and TNF receptor fusion proteins such as those
35 disclosed in USSN 08/406,824 and USSN 08/651,286. In additional, certain virally-encoded proteins are known to bind TNF and act as TNF antagonists, as described in U.S. Patents 5,359,039 and 5,464,938; and inhibitors of an enzyme that cleaves TNF into its biologically active form are also known (see USSN 08/651,363 and USSN 08/655,345).

The relevant disclosures of the aforementioned patents and patent applications are incorporated by reference herein.

IL-17, HVS13 and homologous proteins

- 5 CTLA-8 refers to a cDNA cloned from an activated T cell hybridoma clone (Rouvier et al., *J. Immunol.* 150:5445; 1993). Northern blot analysis indicated that CTLA-8 transcription was very tissue specific. The CTLA-8 gene was found to map at chromosomal site 1a in mice, and at 2q31 in humans. Although a protein encoded by the CTLA-8 gene was never identified by Rouvier et al, the predicted amino acid sequence of
- 10 CTLA-8 was found to be 57% homologous to the predicted amino acid sequence of an ORF present in Herpesvirus Saimiri, HVS13. The CTLA-8 protein is referred to herein as Interleukin-17 (IL-17).

- The complete nucleotide sequence of the genome of HVS has been reported (Albrecht et al., *J. Virol.* 66:5047; 1992). Additional studies on one of the HVS open
- 15 reading frames (ORFs), HVS13, are described in Nicholas et al., *Virol.* 179:1 89; 1990. HVS13 is a late gene which is present in the Hind III-G fragment of HVS. Antisera developed against peptides derived from HVS13 are believed to react with a late protein (Nicholas et al., *supra*).

- As described USSN 08/462,353, a CIP of USSN 08/410,536, filed March 23,
- 20 1995, full length murine CTLA-8 protein and a CTLA-8/Fc fusion protein were expressed, tested, and found to act as a costimulus for the proliferation of T cells. Human IL-17 (CTLA-8) was identified by probing a human T cell library using a DNA fragment derived from degenerate PCR; homologs of IL-17 (CTLA-8) are expected to exist in other species as well. A full length HVS13 protein, as well as an HVS13/Fc fusion protein, were also
- 25 expressed, and found to act in a similar manner to IL-17 (CTLA-8) protein. Moreover, other species of herpesviruses are also likely to encode proteins homologous to that encoded by HVS13.

Proteins and Analogs

- 30 USSN 08/620,694, filed March 21, 1996, discloses isolated IL-17R and homologs thereof having immunoregulatory activity. Such proteins are substantially free of contaminating endogenous materials and, optionally, without associated native-pattern glycosylation. Derivatives of IL-17R within the scope of the invention also include various structural forms of the primary protein which retain biological activity. Due to the presence
- 35 of ionizable amino and carboxyl groups, for example, an IL-17R protein may be in the form of acidic or basic salts, or may be in neutral form. Individual amino acid residues may also be modified by oxidation or reduction.

The primary amino acid structure may be modified by forming covalent or aggregative conjugates with other chemical moieties, such as glycosyl groups, lipids, phosphate, acetyl groups and the like, or by creating amino acid sequence mutants. Covalent derivatives are prepared by linking particular functional groups to amino acid side chains or at the N- or C-termini.

5 Soluble forms of IL-17R are also within the scope of the invention. The nucleotide and predicted amino acid sequence of the murine IL-17R is shown in SEQ ID NOs:1 and 2. Computer analysis indicated that the protein has an N-terminal signal peptide with a cleavage site between amino acid 31 and 32. Those skilled in the art will recognize that the
10 actual cleavage site may be different than that predicted by computer analysis. Thus, the N-terminal amino acid of the cleaved peptide is expected to be within about five amino acids on either side of the predicted cleavage site. The signal peptide is followed by a 291 amino acid extracellular domain, a 21 amino acid transmembrane domain, and a 521 amino acid cytoplasmic tail. Soluble IL-17R comprises the signal peptide and the extracellular domain
15 (residues 1 to 322 of SEQ ID NO:1) or a fragment thereof. Alternatively, a different signal peptide can be substituted for residues 1 through 31 of SEQ ID NO:1.

20 The nucleotide and predicted amino acid sequence of the human IL-17R is shown in SEQ ID NOs:3 and 4. It shares many features with the murine IL-17 R. Computer analysis indicated that the protein has an N-terminal signal peptide with a cleavage site between amino acid 27 and 28. Those skilled in the art will recognize that the actual cleavage site may be different than that predicted by computer analysis. Thus, the N-terminal amino acid of the cleaved peptide is expected to be within about five amino acids on either side of the predicted cleavage site. The signal peptide is followed by a 293 amino acid extracellular domain, a 21 amino acid transmembrane domain, and a 525 amino acid
25 cytoplasmic tail. Soluble IL-17R comprises the signal peptide and the extracellular domain (residues 1 to 320 of SEQ ID NO:1) or a fragment thereof. Alternatively, a different signal peptide can be substituted for the native signal peptide.

Other derivatives of the IL-17R protein and homologs thereof within the scope of this invention include covalent or aggregative conjugates of the protein or its fragments
30 with other proteins or polypeptides, such as by synthesis in recombinant culture as N-terminal or C-terminal fusions. For example, the conjugated peptide may be a signal (or leader) polypeptide sequence at the N-terminal region of the protein which co-translationally or post-translationally directs transfer of the protein from its site of synthesis to its site of function inside or outside of the cell membrane or wall (e.g., the yeast α -factor
35 leader).

Protein fusions can comprise peptides added to facilitate purification or identification of IL-17R proteins and homologs (e.g., poly-His). The amino acid sequence of the inventive proteins can also be linked to an identification peptide such as that

described by Hopp et al., *Bio/Technology* 6:1204 (1988). Such a highly antigenic peptide provides an epitope reversibly bound by a specific monoclonal antibody, enabling rapid assay and facile purification of expressed recombinant protein. The sequence of Hopp et al. is also specifically cleaved by bovine mucosal enterokinase, allowing removal of the peptide from the purified protein. Fusion proteins capped with such peptides may also be resistant to intracellular degradation in *E. coli*.

Soluble forms of some transmembrane proteins have been expressed as fusion proteins in which an extracellular domain of a membrane protein (cognate binding region) is joined to an immunoglobulin heavy chain constant (Fc) domain. Such fusion proteins are useful as reagents to detect their cognate proteins. They are also useful as therapeutic agents in treatment of disease. However, receptors for Fc domains are present on many cell types. Thus, when a fusion protein is formed from an Fc domain and a cognate binding region, binding to a cell may occur either through binding of the cognate binding region to its cognate protein, or through binding of the Fc domain to an Fc receptor (FcR). Such binding of the Fc domain to Fc receptors may overwhelm any binding of the cognate binding region to its cognate. Moreover, binding of Fc domains to Fc receptors induces secretion of various cytokines that are involved in upregulating various aspects of an immune or inflammatory response; such upregulation has been implicated in some of the adverse effects of therapeutic administration of certain antibodies (Krutman et al., *J. Immunol.* 145:1337, 1990; Thistlewaite et al., *Am. J. Kidney Dis.* 11:112, 1988).

Jefferis et al. (*Mol. Immunol.* 27:1237; 1990) reported that a region of an antibody referred to as the hinge region (and specifically residues 234-237 within this region) determine recognition of the antibody by human Fc receptors FcγRI, FcγRII, and FcγRIII. Leu(234) and Leu(235) were critical to high affinity binding of IgG₃ to FcγRI present on U937 cells (Canfield and Morrison, *J. Exp. Med.* 173:1483; 1991). Similar results were obtained by Lund et al. (*J. Immunol.* 147:2657, 1991; *Molecular Immunol.* 29:53, 1991). These authors observed 10-100 fold decrease in affinity of IgG for FcR when a single amino acid substitution was made at a critical residue.

A single amino acid substitution in the Fc domain of an anti-CD3 monoclonal antibody (leucine to glutamic acid at position 235) was found to result in significantly less T cell activation than unmutagenized antibody, while maintaining the immunosuppressive properties (Alegre et al., *J. Immunol.* 148:3461; 1992). Wawrzynczak et al. found that murine monoclonal antibodies that contained a single amino acid substitution at residue 235 had the same serum half-life as did native antibodies (*Mol. Immunol.* 29:221; 1992). Fc domains with reduced affinity for Fc receptors are useful in the preparation of Fc fusion proteins.

Leucine zippers were originally identified in several DNA-binding proteins (Landschulz et al., *Science* 240:1759, 1988). Leucine zipper domain is a term used to refer

to a conserved peptide domain present in these (and other) proteins, which is responsible for dimerization of the proteins. The leucine zipper domain (also referred to herein as an oligomerizing, or oligomer-forming, domain) comprises a repetitive heptad repeat, with four or five leucine residues interspersed with other amino acids. Examples of leucine zipper domains are those found in the yeast transcription factor GCN4 and a heat-stable DNA-binding protein found in rat liver (C/EBP; Landschulz et al., *Science* 243:1681, 1989). Two nuclear transforming proteins, *fos* and *jun*, also exhibit leucine zipper domains, as does the gene product of the murine proto-oncogene, *c-myc* (Landschulz et al., *Science* 240:1759, 1988). The products of the nuclear oncogenes *fos* and *jun* comprise leucine zipper domains preferentially form a heterodimer (O'Shea et al., *Science* 245:646, 1989; Turner and Tjian, *Science* 243:1689, 1989). The leucine zipper domain is necessary for biological activity (DNA binding) in these proteins.

The fusogenic proteins of several different viruses, including paramyxovirus, coronavirus, measles virus and many retroviruses, also possess leucine zipper domains (Buckland and Wild, *Nature* 338:547, 1989; Britton, *Nature* 353:394, 1991; Delwart and Mosialos, *AIDS Research and Human Retroviruses* 6:703, 1990). The leucine zipper domains in these fusogenic viral proteins are near the transmembrane region of the proteins; it has been suggested that the leucine zipper domains could contribute to the oligomeric structure of the fusogenic proteins. Oligomerization of fusogenic viral proteins is involved in fusion pore formation (Spruce et al, *Proc. Natl. Acad. Sci. U.S.A.* 88:3523, 1991). Leucine zipper domains have also been recently reported to play a role in oligomerization of heat-shock transcription factors (Rabindran et al., *Science* 259:230, 1993).

Leucine zipper domains fold as short, parallel coiled coils. (O'Shea et al., *Science* 254:539; 1991) The general architecture of the parallel coiled coil has been well characterized, with a "knobs-into-holes" packing as proposed by Crick in 1953 (*Acta Crystallogr.* 6:689). The dimer formed by a leucine zipper domain is stabilized by the heptad repeat, designated $(abcdefg)_n$ according to the notation of McLachlan and Stewart (*J. Mol. Biol.* 98:293; 1975), in which residues *a* and *d* are generally hydrophobic residues, with *d* being a leucine, which line up on the same face of a helix. Oppositely-charged residues commonly occur at positions *g* and *e*. Thus, in a parallel coiled coil formed from two helical leucine zipper domains, the "knobs" formed by the hydrophobic side chains of the first helix are packed into the "holes" formed between the side chains of the second helix.

The leucine residues at position *d* contribute large hydrophobic stabilization energies, and are important for dimer formation (Krystek et al., *Int. J. Peptide Res.* 38:229, 1991). Lovejoy et al. recently reported the synthesis of a triple-stranded α -helical bundle in which the helices run up-up-down (*Science* 259:1288, 1993). Their studies confirmed that hydrophobic stabilization energy provides the main driving force for the

formation of coiled coils from helical monomers. These studies also indicate that electrostatic interactions contribute to the stoichiometry and geometry of coiled coils.

Several studies have indicated that conservative amino acids may be substituted for individual leucine residues with minimal decrease in the ability to dimerize; multiple changes, however, usually result in loss of this ability (Landschulz et al., *Science* 243:1681, 1989; Turner and Tjian, *Science* 243:1689, 1989; Hu et al., *Science* 250:1400, 1990). van Heekeren et al. reported that a number of different amino residues can be substituted for the leucine residues in the leucine zipper domain of GCN4, and further found that some GCN4 proteins containing two leucine substitutions were weakly active (10 *Nucl. Acids Res.* 20:3721, 1992). Mutation of the first and second heptad leucines of the leucine zipper domain of the measles virus fusion protein (MVF) did not affect syncytium formation (a measure of virally-induced cell fusion); however, mutation of all four leucine residues prevented fusion completely (Buckland et al., *J. Gen. Virol.* 73:1703, 1992). None of the mutations affected the ability of MVF to form a tetramer.

Recently, amino acid substitutions in the *a* and *d* residues of a synthetic peptide representing the GCN4 leucine zipper domain have been found to change the oligomerization properties of the leucine zipper domain (Alber, Sixth Symposium of the Protein Society, San Diego, CA). When all residues at position *a* are changed to isoleucine, the leucine zipper still forms a parallel dimer. When, in addition to this change, (15 all leucine residues at position *d* are also changed to isoleucine, the resultant peptide spontaneously forms a trimeric parallel coiled coil in solution. Substituting all amino acids at position *d* with isoleucine and at position *a* with leucine results in a peptide that tetramerizes. Peptides containing these substitutions are still referred to as leucine zipper domains since the mechanism of oligomer formation is believed to be the same as that for (20 traditional leucine zipper domains such as those described above.

Derivatives of IL-17R may also be used as immunogens, reagents in *in vitro* assays, or as binding agents for affinity purification procedures. Such derivatives may also be obtained by cross-linking agents, such as M-maleimidobenzoyl succinimide ester and N-hydroxysuccinimide, at cysteine and lysine residues. The inventive proteins may also be covalently bound through reactive side groups to various insoluble substrates, such as cyanogen bromide-activated, bisoxirane-activated, carbonyldiimidazole-activated or tosyl-activated agarose structures, or by adsorbing to polyolefin surfaces (with or without glutaraldehyde cross-linking). Once bound to a substrate, proteins may be used to selectively bind (for purposes of assay or purification) antibodies raised against the IL-17R (30 or against other proteins which are similar to the IL-17R, as well as other proteins that bind IL-17R or its homologous proteins.

The present invention also includes IL-17R with or without associated native-pattern glycosylation. Proteins expressed in yeast or mammalian expression systems, e.g.,

COS-7 cells, may be similar or slightly different in molecular weight and glycosylation pattern than the native molecules, depending upon the expression system. Expression of DNAs encoding the inventive proteins in bacteria such as *E. coli* provides non-glycosylated molecules. Functional mutant analogs of IL-17R protein or homologs thereof having inactivated N-glycosylation sites can be produced by oligonucleotide synthesis and ligation or by site-specific mutagenesis techniques. These analog proteins can be produced in a homogeneous, reduced-carbohydrate form in good yield using yeast expression systems. N-glycosylation sites in eukaryotic proteins are characterized by the amino acid triplet Asn-A1-Z, where A1 is any amino acid except Pro, and Z is Ser or Thr. In this sequence, asparagine provides a side chain amino group for covalent attachment of carbohydrate. Such a site can be eliminated by substituting another amino acid for Asn or for residue Z, deleting Asn or Z, or inserting a non-Z amino acid between A1 and Z, or an amino acid other than Asn between Asn and A1.

IL-17R protein derivatives may also be obtained by mutations of the native IL-17R or its subunits. A IL-17R mutated protein, as referred to herein, is a polypeptide homologous to a IL-17R protein but which has an amino acid sequence different from the native IL-17R because of one or a plurality of deletions, insertions or substitutions. The effect of any mutation made in a DNA encoding a IL-17R peptide may be easily determined by analyzing the ability of the mutated IL-17R peptide to inhibit costimulation of T or B cells by IL-17 (CTLA-8) or homologous proteins, or to bind proteins that specifically bind IL-17R (for example, antibodies or proteins encoded by the CTLA-8 cDNA or the HVS13 ORF). Moreover, activity of IL-17R analogs, muteins or derivatives can be determined by any of the assays methods described herein. Similar mutations may be made in homologs of IL-17R, and tested in a similar manner.

Bioequivalent analogs of the inventive proteins may be constructed by, for example, making various substitutions of residues or sequences or deleting terminal or internal residues or sequences not needed for biological activity. For example, cysteine residues can be deleted or replaced with other amino acids to prevent formation of incorrect intramolecular disulfide bridges upon renaturation. Other approaches to mutagenesis involve modification of adjacent dibasic amino acid residues to enhance expression in yeast systems in which KEX2 protease activity is present.

Generally, substitutions should be made conservatively; i.e., the most preferred substitute amino acids are those which do not affect the ability of the inventive proteins to bind their ligands in a manner substantially equivalent to that of native mIL-17R or hIL-17R. Examples of conservative substitutions include substitution of amino acids outside of the binding domain(s), and substitution of amino acids that do not alter the secondary and/or tertiary structure of IL-17R and homologs thereof. Additional examples include substituting one aliphatic residue for another, such as Ile, Val, Leu, or Ala for one another,

or substitutions of one polar residue for another, such as between Lys and Arg; Glu and Asp; or Gln and Asn. Other such conservative substitutions, for example, substitutions of entire regions having similar hydrophobicity characteristics, are well known.

- Similarly, when a deletion or insertion strategy is adopted, the potential effect of the deletion or insertion on biological activity should be considered. Subunits of the inventive proteins may be constructed by deleting terminal or internal residues or sequences. Fragments of IL-17R that bind IL-17 can be readily prepared (for example, by using restriction enzymes to delete portions of the DNA) and tested for their ability to bind IL-17. Additional guidance as to the types of mutations that can be made is provided by a comparison of the sequence of IL-17R to proteins that have similar structures, as well as by performing structural analysis of the inventive proteins.

- Mutations in nucleotide sequences constructed for expression of analog IL-17R must, of course, preserve the reading frame phase of the coding sequences and preferably will not create complementary regions that could hybridize to produce secondary mRNA structures such as loops or hairpins which would adversely affect translation of the receptor mRNA. Although a mutation site may be predetermined, it is not necessary that the nature of the mutation *per se* be predetermined. For example, in order to select for optimum characteristics of mutants at a given site, random mutagenesis may be conducted at the target codon and the expressed mutated viral proteins screened for the desired activity.

- Not all mutations in the nucleotide sequence which encodes a IL-17R protein or homolog thereof will be expressed in the final product, for example, nucleotide substitutions may be made to enhance expression, primarily to avoid secondary structure loops in the transcribed mRNA (see EPA 75,444A, incorporated herein by reference), or to provide codons that are more readily translated by the selected host, e.g., the well-known *E. coli* preference codons for *E. coli* expression.

- Mutations can be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes an analog having the desired amino acid insertion, substitution, or deletion.

- Alternatively, oligonucleotide-directed site-specific mutagenesis procedures can be employed to provide an altered gene having particular codons altered according to the substitution, deletion, or insertion required. Exemplary methods of making the alterations set forth above are disclosed by Walder et al. (*Gene* 42:133, 1986); Bauer et al. (*Gene* 37:73, 1985); Craik (*BioTechniques*, January 1985, 12-19); Smith et al. (*Genetic Engineering: Principles and Methods*, Plenum Press, 1981); and U.S. Patent Nos. 4,518,584 and 4,737,462 disclose suitable techniques, and are incorporated by reference herein.

Due to code degeneracy, there can be considerable variation in nucleotide sequences encoding the same amino acid sequence. Other embodiments include sequences capable of hybridizing under moderately stringent conditions (prewashing solution of 5 X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0) and hybridization conditions of 50°C, 5 X SSC, overnight) to the DNA sequences encoding IL-17R, and other sequences which are degenerate to those which encode the IL-17R. In a preferred embodiment, IL-17R analogs are at least about 70 % identical in amino acid sequence to the amino acid sequence of IL-17R proteins as set forth in SEQ ID NO:1 or SEQ ID NO:3. Similarly, analogs of IL-17R homologs are at least about 70 % identical in amino acid sequence to the amino acid sequence of the native, homologous proteins. In a more preferred embodiment, analogs of IL-17R or homologs thereof are at least about 80 % identical in amino acid sequence to the native form of the inventive proteins; in a most preferred embodiment, analogs of IL-17R or homologs thereof are at least about 90 % identical in amino acid sequence to the native form of the inventive proteins.

Percent identity may be determined using a computer program, for example, the GAP computer program described by Devereux et al. (*Nucl. Acids Res.* 12:387, 1984) and available from the University of Wisconsin Genetics Computer Group (UWGCG). For fragments derived from the IL-17R protein, the identity is calculated based on that portion of the IL-17R protein that is present in the fragment. Similar methods can be used to analyze homologs of IL-17R.

The ability of IL-17R analogs to bind CTLA-8 can be determined by testing the ability of the analogs to inhibit IL-17 (CTLA-8) -induced T cell proliferation. Alternatively, suitable assays, for example, an enzyme immunoassay or a dot blot, employing CTLA-8 or HSV13 (or a homolog thereof which binds native IL-17R) can be used to assess the ability of IL-17R analogs to bind CTLA-8. Such methods are well known in the art.

Expression of Recombinant Receptors for IL-17

The proteins of the present invention are preferably produced by recombinant DNA methods by inserting a DNA sequence encoding IL-17R protein or a homolog thereof into a recombinant expression vector and expressing the DNA sequence in a recombinant microbial expression system under conditions promoting expression. DNA sequences encoding the proteins provided by this invention can be assembled from cDNA fragments and short oligonucleotide linkers, or from a series of oligonucleotides, to provide a synthetic gene which is capable of being inserted in a recombinant expression vector and expressed in a recombinant transcriptional unit.

Recombinant expression vectors include synthetic or cDNA-derived DNA fragments encoding IL-17R, homologs, or bioequivalent analogs, operably linked to suitable transcriptional or translational regulatory elements derived from mammalian,

microbial, viral or insect genes. Such regulatory elements include a transcriptional promoter, an optional operator sequence to control transcription, a sequence encoding suitable mRNA ribosomal binding sites, and sequences which control the termination of transcription and translation, as described in detail below. The ability to replicate in a host, usually conferred by an origin of replication, and a selection gene to facilitate recognition of transformants may additionally be incorporated.

DNA regions are operably linked when they are functionally related to each other. For example, DNA for a signal peptide (secretory leader) is operably linked to DNA for a polypeptide if it is expressed as a precursor which participates in the secretion of the polypeptide; a promoter is operably linked to a coding sequence if it controls the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to permit translation. Generally, operably linked means contiguous and, in the case of secretory leaders, contiguous and in reading frame. DNA sequences encoding IL-17R or homologs which are to be expressed in a microorganism will preferably contain no introns that could prematurely terminate transcription of DNA into mRNA.

Useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed. *E. coli* is typically transformed using derivatives of pBR322, a plasmid derived from an *E. coli* species (Bolivar et al., *Gene* 2:95, 1977). pBR322 contains genes for ampicillin and tetracycline resistance and thus provides simple means for identifying transformed cells.

Promoters commonly used in recombinant microbial expression vectors include the β -lactamase (penicillinase) and lactose promoter system (Chang et al., *Nature* 275:615, 1978; and Goeddel et al., *Nature* 281:544, 1979), the tryptophan (trp) promoter system (Goeddel et al., *Nucl. Acids Res.* 8:4057, 1980; and EPA 36,776) and tac promoter (Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, p. 412, 1982). A particularly useful bacterial expression system employs the phage λ PL promoter and cI857ts thermolabile repressor. Plasmid vectors available from the American Type Culture Collection which incorporate derivatives of the λ PL promoter include plasmid pHUB2, resident in *E. coli* strain JMB9 (ATCC 37092) and pPLc28, resident in *E. coli* RR1 (ATCC 53082).

Suitable promoter sequences in yeast vectors include the promoters for metallothionein, 3-phosphoglycerate kinase (Hitzeman et al., *J. Biol. Chem.* 255:2073,

1980) or other glycolytic enzymes (Hess et al., *J. Adv. Enzyme Reg.* 7:149, 1968; and Holland et al., *Biochem.* 17:4900, 1978), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. Suitable vectors and promoters
5 for use in yeast expression are further described in R. Hitzeman et al., EPA 73,657.

Preferred yeast vectors can be assembled using DNA sequences from pBR322 for selection and replication in *E. coli* (Amp^r gene and origin of replication) and yeast DNA sequences including a glucose-repressible ADH2 promoter and α -factor secretion leader.
10 The ADH2 promoter has been described by Russell et al. (*J. Biol. Chem.* 258:2674, 1982) and Beier et al. (*Nature* 300:724, 1982). The yeast α -factor leader, which directs secretion of heterologous proteins, can be inserted between the promoter and the structural gene to be expressed. See, e.g., Kurjan et al., *Cell* 30:933, 1982; and Bitter et al., *Proc. Natl. Acad. Sci. USA* 81:5330, 1984. The leader sequence may be modified to contain, near its 3'
15 end, one or more useful restriction sites to facilitate fusion of the leader sequence to foreign genes.

The transcriptional and translational control sequences in expression vectors to be used in transforming vertebrate cells may be provided by viral sources. For example, commonly used promoters and enhancers are derived from Polyoma, Adenovirus 2,
20 Simian Virus 40 (SV40), and human cytomegalovirus. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites may be used to provide the other genetic elements required for expression of a heterologous DNA sequence. The early and late promoters are particularly useful because both are obtained easily from the virus as a fragment which also contains the
25 SV40 viral origin of replication (Fiers et al., *Nature* 273:113, 1978). Smaller or larger SV40 fragments may also be used, provided the approximately 250 bp sequence extending from the *Hind* III site toward the *Bgl* II site located in the viral origin of replication is included. Further, viral genomic promoter, control and/or signal sequences may be utilized, provided such control sequences are compatible with the host cell chosen.
30 Exemplary vectors can be constructed as disclosed by Okayama and Berg (*Mol. Cell. Biol.* 3:280, 1983).

A useful system for stable high level expression of mammalian receptor cDNAs in C127 murine mammary epithelial cells can be constructed substantially as described by Cosman et al. (*Mol. Immunol.* 23:935, 1986). A preferred eukaryotic vector for
35 expression of IL-17R DNA is referred to as pDC406 (McMahan et al., *EMBO J.* 10:2821, 1991), and includes regulatory sequences derived from SV40, human immunodeficiency virus (HIV), and Epstein-Barr virus (EBV). Other preferred vectors include pDC409 and pDC410, which are derived from pDC406. pDC410 was derived from pDC406 by

substituting the EBV origin of replication with sequences encoding the SV40 large T antigen. pDC409 differs from pDC406 in that a *Bgl* II restriction site outside of the multiple cloning site has been deleted, making the *Bgl* II site within the multiple cloning site unique.

- 5 A useful cell line that allows for episomal replication of expression vectors, such as pDC406 and pDC409, which contain the EBV origin of replication, is CV-1/EBNA (ATCC CRL 10478). The CV-1/EBNA cell line was derived by transfection of the CV-1 cell line with a gene encoding Epstein-Barr virus nuclear antigen-1 (EBNA-1) and constitutively express EBNA-1 driven from human CMV immediate-early enhancer/promoter.

10

Host Cells

- Transformed host cells are cells which have been transformed or transfected with expression vectors constructed using recombinant DNA techniques and which contain sequences encoding the proteins of the present invention. Transformed host cells may
15 express the desired protein (IL-17R or homologs thereof), but host cells transformed for purposes of cloning or amplifying the inventive DNA do not need to express the protein. Expressed proteins will preferably be secreted into the culture supernatant, depending on the DNA selected, but may be deposited in the cell membrane.

- Suitable host cells for expression of viral proteins include prokaryotes, yeast or
20 higher eukaryotic cells under the control of appropriate promoters. Prokaryotes include gram negative or gram positive organisms, for example *E. coli* or *Bacillus* spp. Higher eukaryotic cells include established cell lines of mammalian origin as described below. Cell-free translation systems could also be employed to produce viral proteins using RNAs derived from the DNA constructs disclosed herein. Appropriate cloning and expression
25 vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described by Pouwels et al. (*Cloning Vectors: A Laboratory Manual*, Elsevier, New York, 1985), the relevant disclosure of which is hereby incorporated by reference.

- Prokaryotic expression hosts may be used for expression of IL-17R or homologs that do not require extensive proteolytic and disulfide processing. Prokaryotic expression
30 vectors generally comprise one or more phenotypic selectable markers, for example a gene encoding proteins conferring antibiotic resistance or supplying an autotrophic requirement, and an origin of replication recognized by the host to ensure amplification within the host. Suitable prokaryotic hosts for transformation include *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and various species within the genera *Pseudomonas*, *Streptomyces*, and
35 *Staphylococcus*, although others may also be employed as a matter of choice.

Recombinant IL-17R may also be expressed in yeast hosts, preferably from the *Saccharomyces* species, such as *S. cerevisiae*. Yeast of other genera, such as *Pichia* or *Kluyveromyces* may also be employed. Yeast vectors will generally contain an origin of

- replication from the 2 μ yeast plasmid or an autonomously replicating sequence (ARS), promoter, DNA encoding the viral protein, sequences for polyadenylation and transcription termination and a selection gene. Preferably, yeast vectors will include an origin of replication and selectable marker permitting transformation of both yeast and *E. coli*, e.g., the ampicillin resistance gene of *E. coli* and *S. cerevisiae* *trp1* gene, which provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, and a promoter derived from a highly expressed yeast gene to induce transcription of a structural sequence downstream. The presence of the *trp1* lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan.

- Suitable yeast transformation protocols are known to those of skill in the art; an exemplary technique is described by Hinnen et al., *Proc. Natl. Acad. Sci. USA* 75:1929, 1978, selecting for Trp⁺ transformants in a selective medium consisting of 0.67% yeast nitrogen base, 0.5% casamino acids, 2% glucose, 10 μ g/ml adenine and 20 μ g/ml uracil. Host strains transformed by vectors comprising the ADH2 promoter may be grown for expression in a rich medium consisting of 1% yeast extract, 2% peptone, and 1% glucose supplemented with 80 μ g/ml adenine and 80 μ g/ml uracil. Derepression of the ADH2 promoter occurs upon exhaustion of medium glucose. Crude yeast supernatants are harvested by filtration and held at 4°C prior to further purification.

- Various mammalian or insect cell culture systems can be employed to express recombinant protein. Baculovirus systems for production of heterologous proteins in insect cells are reviewed by Luckow and Summers, *BioTechnology* 6:47 (1988). Examples of suitable mammalian host cell lines include the COS-7 lines of monkey kidney cells, described by Gluzman (*Cell* 23:175, 1981), and other cell lines capable of expressing an appropriate vector including, for example, CV-1/EBNA (ATCC CRL 10478), L cells, C127, 3T3, Chinese hamster ovary (CHO), HeLa and BHK cell lines. Mammalian expression vectors may comprise nontranscribed elements such as an origin of replication, a suitable promoter and enhancer linked to the gene to be expressed, and other 5' or 3' flanking nontranscribed sequences, and 5' or 3' nontranslated sequences, such as necessary ribosome binding sites, a polyadenylation site, splice donor and acceptor sites, and transcriptional termination sequences.

Purification of Receptors for IL-17

- Purified IL-17R, homologs, or analogs are prepared by culturing suitable host/vector systems to express the recombinant translation products of the DNAs of the present invention, which are then purified from culture media or cell extracts. For example, supernatants from systems which secrete recombinant protein into culture media

can be first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit.

Following the concentration step, the concentrate can be applied to a suitable purification matrix. For example, a suitable affinity matrix can comprise a counter structure protein or lectin or antibody molecule bound to a suitable support. Alternatively, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. Sulfopropyl groups are preferred. Gel filtration chromatography also provides a means of purifying the inventive proteins.

Affinity chromatography is a particularly preferred method of purifying IL-17R and homologs thereof. For example, a IL-17R expressed as a fusion protein comprising an immunoglobulin Fc region can be purified using Protein A or Protein G affinity chromatography. Moreover, a IL-17R protein comprising an oligomerizing zipper domain may be purified on a resin comprising an antibody specific to the oligomerizing zipper domain. Monoclonal antibodies against the IL-17R protein may also be useful in affinity chromatography purification, by utilizing methods that are well-known in the art. A ligand (i.e., IL-17 or HVS-13) may also be used to prepare an affinity matrix for affinity purification of IL-17R.

Finally, one or more reversed-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify a IL-17R composition. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a homogeneous recombinant protein.

Recombinant protein produced in bacterial culture is usually isolated by initial extraction from cell pellets, followed by one or more concentration, salting-out, aqueous ion exchange or size exclusion chromatography steps. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps. Microbial cells employed in expression of recombinant viral protein can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Fermentation of yeast which express the inventive protein as a secreted protein greatly simplifies purification. Secreted recombinant protein resulting from a large-scale fermentation can be purified by methods analogous to those disclosed by Urdal et al. (*J. Chromatog.* 296:171, 1984). This reference describes two sequential, reversed-phase

HPLC steps for purification of recombinant human GM-CSF on a preparative HPLC column.

- Protein synthesized in recombinant culture is characterized by the presence of cell components, including proteins, in amounts and of a character which depend upon the purification steps taken to recover the inventive protein from the culture. These components ordinarily will be of yeast, prokaryotic or non-human higher eukaryotic origin and preferably are present in innocuous contaminant quantities, on the order of less than about 1 percent by weight. Further, recombinant cell culture enables the production of the inventive proteins free of other proteins which may be normally associated with the proteins as they are found in nature in the species of origin.

Administration of IL-17R Compositions

- The present invention provides methods of using therapeutic compositions comprising an effective amount of a protein and a suitable diluent and carrier. The use of IL-17R or homologs in conjunction with soluble cytokine receptors or cytokines, or other immunoregulatory molecules is also contemplated. Such molecules can be administered separately, sequentially or simultaneously with IL-17R compositions. Particularly preferred immunoregulatory molecules are soluble IL-1 receptors, soluble TNF receptors, and fusion proteins thereof.

- For therapeutic use, purified protein is administered to a patient, preferably a human, for treatment in a manner appropriate to the indication. Thus, for example, IL-17R protein compositions administered to regulate NO levels can be given by bolus injection, continuous infusion, sustained release from implants, or other suitable technique. Typically, a therapeutic agent will be administered in the form of a composition comprising purified IL-17R, in conjunction with physiologically acceptable carriers, excipients or diluents. Such carriers will be nontoxic to recipients at the dosages and concentrations employed.

- Ordinarily, the preparation of such protein compositions entails combining the inventive protein with buffers, antioxidants such as ascorbic acid, low molecular weight (less than about 10 residues) polypeptides, proteins, amino acids, carbohydrates including glucose, sucrose or dextrans, chelating agents such as EDTA, glutathione and other stabilizers and excipients. Neutral buffered saline or saline mixed with conspecific serum albumin are exemplary appropriate diluents. Preferably, product is formulated as a lyophilizate using appropriate excipient solutions (e.g., sucrose) as diluents. Appropriate dosages can be determined in trials. The amount and frequency of administration will depend, of course, on such factors as the nature and severity of the indication being treated, the desired response, the condition of the patient, and so forth.

Receptors for IL-17 (CTLA-8) can be administered for the purpose of regulating levels of NO. Soluble IL-17R are thus likely to be useful in treatment of osteoarthritis. The inventive receptor proteins will also be useful for prevention or treatment inflammation.

The following examples are offered by way of illustration, and not by way of limitation. Those skilled in the art will recognize that variations of the invention embodied in the examples can be made, especially in light of the teachings of the various references cited herein, the disclosures of which are incorporated by reference.

EXAMPLE 1

This example illustrates the ability of IL-17R to inhibit the proliferative response of T cells to mitogens. Lymphoid organs were harvested aseptically and cell suspension was created. Splenic and lymph node T cells were isolated from the cell suspension. The purity of the resulting splenic T cell preparations was routinely >95% CD3⁺ and <1% sIgM⁺. Purified murine splenic T cells (2x10⁵/well) were cultured with either 1% PHA or 1 µg/ml Con A, and a soluble IL-17R (a soluble form of IL-17R comprising the extracellular region of IL-17R fused to the Fc region of human IgG1) was titrated into the assay. Proliferation was determined after 3 days with the addition of 1 µCi [³H]thymidine.

Secretion of cytokines (Interleukin-2) was determined for murine T cells cultured for 24 hr with 1 µg/ml of Con A in the presence or absence of 10 µg/ml of IL-17R.Fc or in the presence of a control Fc protein. IL-2 production was measured by ELISA and results expressed as ng/ml IL-2 produced.

Soluble IL-17R/Fc significantly inhibited the mitogen-induced proliferation of purified murine splenic T cells in a dose dependent manner, while a control Fc had no effect on the murine T cell proliferation. Complete inhibition of mitogen induced proliferation was observed at a soluble IL-17R.Fc concentration of 10 µg/ml. Analysis of IL-2 production by splenic T cells activated with Con A in the presence or absence of IL-17R.Fc in the culture revealed that addition of IL-17R.Fc to the T-cell culture inhibited IL-2 production to levels 8-9-fold lower than those observed in cultures containing media alone or media plus a control Fc protein. Similar results were observed when purified human T cells were used.

EXAMPLE 2

This example illustrates the ability of IL-17R to inhibit the production of NO by cartilage-associated cells. Articular cartilage is obtained from OA-affected patients or normal controls substantially as described in Amin et al., supra. The cartilage is cut into small (approximately 3 mm) discs, which are placed in organ culture in the presence or

- absence of IL-17R.Fc or in the presence of a control Fc protein. Nitric oxide production is assayed by determining the nitrite level in the medium at different time intervals, for example by using a modified Griess reaction (*Anal. Biochem.* 12b:12299; 1982). Ding et al. (*J. Immunol.* 141:2407, 1988) also describe a useful method of measuring NO in *ex vivo* organ cultures of synovium and cartilage associated cells. The IL-17R.Fc is titrated to determine an effective concentration to inhibit NO production. Other soluble forms of IL-17R are also used to regulate NO levels in this manner.
- 5

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2